EGOZITE CRADLE CAP LOTION

Salicylic Acid (6% w/w)

What is in this leaflet

- 1. What Egozite Cradle Cap Lotion is used for
- 2. How Egozite Cradle Cap Lotion works
- 3. Before you use Egozite Cradle Cap Lotion
- 4. How to use Egozite Cradle Cap Lotion
- 5. While you are using it
- 6. Side effects
- 7. Storage and disposal of Egozite Cradle Cap Lotion
- 8. Product description
- 9. Manufacturer and Product Registration Holder
- 10. Date of revision

What Egozite Cradle Cap Lotion is used for

- Especially formulated to treat cradle cap (a yellowish, patchy, greasy, scaly and crusty skin rash that occurs on the scalp of recently born babies)
- Helps loosen the crusts and remove the cradle cap

How Egozite Cradle Cap Lotion works

- Egozite Cradle Cap Lotion contains salicylic acid for loosening scales & softening

Before you use Egozite Cradle Cap Lotion

- When you must not use it

This product should not be used by individuals with extreme inflammation of the scalp, weeping eczema (inflammatory form of skin disease marked by fluid exudate) or known sensitivity to any ingredients of this product.

- Before you start to use it
Please discuss with your doctor or
pharmacist before using the
product.

For use during breastfeeding, no deleterious effects are expected as the ingredients have been widely used over many years and no adverse effects have been reported in published literature. - *Taking other medicines*Currently there have been no reports of problems when this

product is used with other

How to use Egozite Cradle Cap Lotion

- How much to use

medicines.

Apply to crusts only avoiding noncrusted areas. Wipe excess lotion from non-crusted areas with dry cotton wool.

- When to use it

Apply twice daily for 3-5 days without washing hair, then wash with a gentle shampoo.

- How long to use it

Continue using Egozite Cradle Cap Lotion on any remaining crusts twice daily. Please follow the advice of the doctor or pharmacist.

- If you forget to use it

If forget to use this medicine, use it as soon as remember. Then next, use this medicine at the usual time.

- If you use too much (overdose)

No information is available concerning overdosage in human beings following topical application.

While you are using it

- Things you must do

Avoid contact with eyes. For external use only. Ensure scalp is dry. Avoid soap.

- Things you must not do

Do not swallow. Do not apply to red and non-crusted areas. Do not remove crusts forcibly.

- Things to be careful of

If irritation occurs, discontinue use.

Side effects

Allergy to any ingredient may cause skin irritation.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website npra.moh.gov.my (Public → Reporting Medicinal Problems / Side Effects / AEFI / Vaccine Safety).

Storage and Disposal of Egozite
Cradle Cap Lotion

- Storage

Store below 30°C. Keep out of reach of children.

- Disposal

Egozite Cradle Cap Lotion should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of Egozite Cradle Cap Lotion that is no longer required. These measures will help to protect the environment.

Product Description

- What it looks like

Egozite Cradle Cap Lotion is a clear oily, yellow, viscous liquid in 50 mL bottles.

- Ingredients

- Active ingredients: salicylic acid.
- Inactive ingredients: olive oil, propyl gallate, castor oil.

- MAL number:

MAL19912422XRZ

Manufacturer

Ego Pharmaceuticals Pty Ltd 21-31 Malcolm Road, Braeside, 3195 Australia

Product Registration Holder

Ego Pharm Malaysia Sdn Bhd 1-6 Block A, 72A Jln Universiti, 46100 Petaling Jaya, Selangor Darul Ehsan.

EGOZITE CRADLE CAP LOTION

Salicylic Acid (6% w/w)

Date of revision 21/08/2017

Serial Number NPRA(R4/1)140817/00221