

# PHARMANIAGA PSEUDOEPHEDRINE T TABLET

Pseudoephedrine hydrochloride/Triprolidine hydrochloride (60mg/2.5mg per 5ml)

## What is in this leaflet

1. What Pharmaniaga Pseudoephedrine T Tablet is used for
2. How Pharmaniaga Pseudoephedrine T Tablet works
3. Before you use Pharmaniaga Pseudoephedrine T Tablet
4. How to use Pharmaniaga Pseudoephedrine T Tablet
5. While you are using Pharmaniaga Pseudoephedrine T Tablet
6. Side effects
7. Storage and Disposal of Pharmaniaga Pseudoephedrine T Tablet
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of Revision

## What Pharmaniaga

### Pseudoephedrine T Tablet is used for

Pseudoephedrine T Tablet is a medicine which is used to relieve the symptoms of colds, flu and allergies including sneezing, runny nose, watery eyes, nasal congestion and blocked sinuses. The tablet contains pseudoephedrine hydrochloride, which is a decongestant that relieves nasal and sinus congestion and triprolidine hydrochloride which is an antihistamine that helps stop sneezing, runny nose and watery eyes.

## How Pharmaniaga

### Pseudoephedrine T Tablet works

*Triprolidine* works by lowering the level of a naturally-occurring chemical in the body called histamine, which triggers the reaction that leads to allergy symptoms.

*Pseudoephedrine* works by causing the blood vessels in the sinuses to tighten, decreasing the swelling of the sinuses and making it easier to breathe.

## Before you use Pharmaniaga

### Pseudoephedrine T Tablet

#### *- When you must not use it*

Do not take Pharmaniaga Pseudoephedrine T Tablet if you:

- are allergic to any medicine containing pseudoephedrine, triprolidine or any of the ingredients listed below
- have an overactive thyroid gland.
- have glaucoma (increased pressure in the eye)
- have prostate problems (difficulty with passing water or needing to pass water often)
- very high blood pressure or severe heart disease
- are taking, or have taken in the last two weeks, medicines for depression known as Monoamine Oxidase Inhibitors (MAOIs).
- are taking other sympathomimetic medicines (*stimulants or appetite suppressants* and medicines used to treat congestion and asthma)

#### *- Before you start to use it*

Tell your pharmacist if you have allergies to any other medicines, foods, preservatives or dyes.

Tell your doctor or pharmacist if you are pregnant or plan to become pregnant or are breast-feeding. Your doctor or pharmacist can discuss with you the risks and benefits involved.

Talk to your doctor or pharmacist if you:

- have high blood pressure or heart disease.
- have diabetes or if you been told by your doctor that you have an intolerance to some sugars.
- are taking alcohol.
- have epilepsy.

#### *Paediatric use*

Use is not recommended in newborn or premature infants and in children under 6 years.

## *Elderly use*

Confusion, dizziness, sedation, low blood pressure, hyperexcitability and anti-cholinergic side-effects, such as dryness of mouth and urinary retention (especially in males) may be more likely to occur in elderly people taking anti-histamines. In addition, elderly patients are also more sensitive to the blood vessel contraction and central nervous system effect.

## *Pregnancy and breastfeeding*

Do not take Pharmaniaga Pseudoephedrine T Tablet if you are pregnant, trying to get pregnant or think you may be pregnant.

## *Do not take Pharmaniaga*

Pseudoephedrine T Tablet if you are breast-feeding. Small amounts of the medicine may be excreted in breast milk. Ask your doctor or pharmacist for advice before taking any medicine.

## *- Taking other medicines*

Tell your pharmacist if you are taking any other medicines, including any that you get without a prescription from your pharmacy, supermarket or health food shop.

Tell your doctor or pharmacist if you are taking any other medicines, including:

- *Sedatives* (drugs used to treat anxiety and tension).
- *Antihypertensives or diuretics* (drugs used to treat high blood pressure such as *methyldopa, alpha and beta blockers*)
- *Tricyclic antidepressants* (used to treat mood disorders).
- *Halogenated anaesthetic agents* (such as *chloroform, cyclopropane, halothane, enflurane or isoflurane*).
- *Sympathomimetic agent* (stimulants or appetite suppressants and drugs used to treat congestion and asthma)

# PHARMANIAGA PSEUDOEPHEDRINE T TABLET

Pseudoephedrine hydrochloride/Triprolidine hydrochloride (60mg/2.5mg per 5ml)

- *Anticholinergics agent such as atropine.*
- *Monoamine oxidase inhibitors (MAOIs) (used to treat depression).*

## **How to use Pharmaniaga Pseudoephedrine T Tablet**

Follow all directions given to you by your pharmacist or doctor carefully. They may differ from the information contained in this leaflet. If you do not understand the directions on your prescription bottle, ask your pharmacist, nurse, or doctor to explain the instructions to you.

### **-How much to use**

For oral use only.

#### **Adults :**

One tablet to be taken 3 times daily.

#### **Children 6-12 years:**

Half a tablet to be taken 3 times daily.

Not suitable to be used for children below 2 years of age. Use carefully and follow the advice from doctor or pharmacist for children aged 2 to 6 years.

### **-When to use it**

Leave at least 4 hours between doses.

- Do not give more than 4 doses in 24 hours.
- Use only when simple measures have failed to provide adequate relief.

### **-How long to use it**

Do not use for more than 5 days without consulting your doctor or pharmacist.

This medicine is usually only used for short periods of time. If symptoms persist, consult your doctor.

### **-If you forget to use it**

If you forget to take a dose, take the next dose when needed provided that the last dose was taken at least 4 hours ago.

Do not take a double dose.

### **-If you use too much (overdose)**

Contact your doctor or go to your nearest hospital immediately. Do not drive if you have taken too much. Symptoms of overdoses include:

- Anti-cholinergic effects: Clumsiness or unsteadiness, severe dryness of mouth, nose or throat, flushing or redness of face, shortness of breath or troubled breathing.
- CNS stimulation: Hallucinations, seizures, trouble in sleeping.
- Severe drowsiness.
- High blood pressure: Continuing headache, slow or fast heartbeat.

### **While you are using it**

#### **-Things you must do**

Talk to your pharmacist or doctor if your symptoms do not improve. Your pharmacist or doctor will assess your condition and decide if you should continue to take the medicine.

Elderly people should inform their doctor they are taking this medicine as their doctor may wish to monitor their kidney and liver function.

#### **-Things you must not do**

Do not take Pharmaniaga Pseudoephedrine T Tablet to treat any other complaints unless your pharmacist or doctor tells you to.

Do not give your medicine to anyone else, even if they have the same condition as you.

Do not take more than the recommended dose unless your pharmacist or doctor tells you to

### **-Things to be careful of**

This medicine may cause drowsiness. If affected, do not drive or operate machinery. Avoid alcoholic drink.

### **Side effects**

Tell your pharmacist or doctor if you do not feel well while you are taking Pharmaniaga Pseudoephedrine T Tablet. This medicine helps most people, but it may have unwanted side effects in a few people. All medicines can have side effects. Sometimes they are serious, most of the time they are not. You may need medical attention if you get some of the side effects.

Do not be alarmed by the following list of side effects. You may not experience any of them.

Ask your pharmacist/doctor to answer any questions you may have. Get emergency medical help if you have any of these signs of an allergic reaction while taking Pharmaniaga Pseudoephedrine T Tablet: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Stop using this medication and call your doctor at once if you have a serious side effect such as:

- fast or uneven heart rate;
- mood changes;
- tremor, seizure (convulsions);
- easy bruising or bleeding, unusual weakness;
- urinating less than usual or not at all;
- feeling short of breath; or
- dangerously high blood pressure (severe headache, blurred vision, buzzing in your ears, anxiety, chest pain, uneven heartbeats, seizure).

Less serious side effects of *pseudoephedrine/triprolidine* may include:

- dizziness, drowsiness;
- dry mouth, nose, or throat;
- blurred vision; or

# PHARMANIAGA PSEUDOEPHEDRINE T TABLET

Pseudoephedrine hydrochloride/Triprolidine hydrochloride (60mg/2.5mg per 5ml)

- feeling nervous or restless.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website [portal.bpfk.gov.my](http://portal.bpfk.gov.my)(Consumers → Reporting)

## **Storage and Disposal of Pharmaniaga Pseudoephedrine T Tablet**

### **-Storage**

Keep the medicines in a cool dry place where the temperature stays below 25°C. Protect from light.

Do not store it or any other medicine in the bathroom, or near a sink, or on a window sill.

Do not leave it in the car on hot days.

Heat and dampness can destroy some medicines.

Do not use this medication after the expiry date which is printed on the product packaging. The expiry date refers to the last day of that month. Keep it where children cannot reach it.

### **-Disposal**

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

## **Product Description**

### **-What it looks like**

Yellow, ellipse shaped tablet with Pharmaniaga icon on one side and scoreline on the other.

### **-Ingredients:**

#### **-Active ingredient(s):**

Pseudoephedrine T Tablet contains Pseudoephedrine HCl 60mg, Triprolidine HCl 2.5mg

#### **-Inactive ingredient(s):**

Ac-Di-Sol, Kollidon, Quinoline Yellow Lake, Lactose, Aerosil, Cornstarch, Isopropyl Alcohol and Magnesium Stearate.

### **-MAL Number:**

Pharmaniaga Pseudoephedrine T Tablet: MAL19971856A

### **Manufacturer:**

Pharmaniaga Manufacturing Berhad (60016-D),  
11A, Jalan P/1, Kawasan Perusahaan Bangi, 43650 Bandar Baru Bangi, Selangor Darul Ehsan, Malaysia.

### **Product Registration Holder:**

Pharmaniaga Manufacturing Berhad (60016-D),  
11A, Jalan P/1, Kawasan Perusahaan Bangi, 43650 Bandar Baru Bangi, Selangor Darul Ehsan, Malaysia.

### **Date of revision:**

28/10/2015

### **Serial number**

BPFK(R4/1)271015/00382