

CLAMENTIN FILM COATED TABLET

Amoxicillin /Clavulanic Acid (375mg, 625mg)

What is in this leaflet

1. What CLAMENTIN F.C TABLET is used for
2. How CLAMENTIN F.C TABLET works
3. Before you use CLAMENTIN F.C TABLET
4. How to use CLAMENTIN F.C TABLET
5. While you are using CLAMENTIN F.C TABLET
6. Side effects
7. Storage and disposal of CLAMENTIN F.C TABLET
8. Product description
9. Manufacturer and Product Registration Holder
10. Date of revision

What CLAMENTIN F.C TABLET is used for

It is used for short-term treatment of bacterial infection at respiratory tract, urinary tract, skin and soft tissue, bone and joint and dental.

Your doctor may have prescribed CLAMENTIN F.C TABLET for another reason. Ask your doctor if you have any questions about why CLAMENTIN F.C TABLET has been prescribed for you.

How CLAMENTIN F.C TABLET works

CLAMENTIN F.C TABLET consists of active substance Amoxicillin/Clavulanic Acid. These medicine work by killing the bacteria, which cause infections.

Before you use CLAMENTIN F.C TABLET

-When you must not use it

Do not use CLAMENTIN F.C TABLET if:

-you have had an allergic reaction to penicillin or similar types of antibiotics (such as cephalosporins) or any of the ingredients contained in CLAMENTIN F.C TABLET
The ingredients are listed at the end of this leaflet. Some of the symptoms of an allergic reactions may include skin rash, itching, difficulty in breathing and swelling of the face or tongue.

-you have previously experienced liver problems after using CLAMENTIN F.C TABLET or any other penicillin antibiotic.

-Before you start to use it

Tell your doctor if:

-you have ever had an allergic reaction (such as a rash) to antibiotics such as penicillin or other substances in the past.

-you are allergic to foods, dyes, preservatives

or any other medicines.

-you have experienced liver problems after using CLAMENTIN F.C TABLET or any other medicines.

-you have glandular fever (mononucleosis).

-you have kidney or liver problem.

Pregnancy and lactation

Please consult your doctor before start to use CLAMENTIN F.C TABLET if you are pregnant, plan to become pregnant, breast-feeding or intend to breast-feed. Your doctor will have weighed the risks of you taking CLAMENTIN F.C TABLET against the benefits they expect it will have for you.

CLAMENTIN F.C TABLET 625mg is not recommended for children of 12 years and under.

-Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

Some medicines when taken with CLAMENTIN F.C TABLET may interfere with each other. These include:

-anticoagulant therapy, medicines for blood thinning effect.

-oral contraceptives.

-probenecid, medicine for gout.

-allopurinol, medicine for gout.

Your doctor or pharmacist may have more information on medicines to be careful with or to avoid while taking medication.

How to use CLAMENTIN F.C TABLET

-How much to use

Adults and children over 12 years

Mild – Moderate infections

: One tablet CLAMENTIN F.C TABLET 625mg two times a day.

: One tablet CLAMENTIN F.C TABLET 375mg three times a day

Your doctor might prescribes a lower dosage for you, if you have kidney or liver disease.

Follow all directions given to you by your doctor and pharmacist carefully. They may differ from the information contained in this leaflet. If you do not

understand the instructions on the label, ask your doctor or pharmacist for help.

-When to use it

Use as directed by your doctor or pharmacist

-How long to use it

Continue using CLAMENTIN F.C TABLET for as long as your doctor recommends.

Do not stop taking medication, even if you feel better after a few days, unless advised by your doctor. Your infection may not clear completely if you stop taking your medication too soon.

-If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it.

If you miss a dose of CLAMENTIN F.C TABLET and you are taking it regularly, take it as soon as possible. If it is approaching time for the next dose, do not double the dose to make up for the forgotten dose, unless advised by your health care provider. Do not take 2 doses at once.

-If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have use too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Taking too many tablets may cause gastrointestinal symptoms and disturbance of the fluid and electrolytes balance.

While you are using CLAMENTIN F.C TABLET

-Things you must do

Follow your doctor's instruction when you are taking CLAMENTIN F.C TABLET.

Tell all the doctors, dentists and pharmacists treating you that you are taking CLAMENTIN F.C TABLET.

Tell your doctor immediately if you become pregnant and intended to breast-feed while taking this medicine.

Ask your doctor or pharmacist for advice before taking any medicine.

If you get severe diarrhoea, tell your doctor or pharmacist immediately.

-Things you must not do

Do not stop taking this medicine or change the dosing regimen without instruction from your doctor. Do not give CLAMENTIN F.C TABLET to anyone else, even if they have the same condition as you.

-Things to be careful of

Driving and using machine

Be careful driving or operating machinery until you know how CLAMENTIN F.C TABLET affects you.

Side effects

CLAMENTIN F.C TABLET may cause gastrointestinal side effect such as diarrhea, indigestion, nausea and vomiting. It may give hematological (blood disorder) effects such as thrombocytopenia, haemolytic anemia and cause hypersensitivity reactions such as urticarial (itching) and erythematous rashes, severe allergic reaction such as Stevens-Johnson syndrome, toxic epidermal necrolysis and exfoliative dermatitis. It also could cause hyperactivity, dizziness, headache and phlebitis (inflammation of vein).

Like all medicines, CLAMENTIN F.C TABLET can cause side effects, although not everybody gets them.

Visit your doctor or pharmacist immediately if you are experience any other side effects after taking this medicine.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website portal.bpfk.gov.my (Consumers → Reporting)

Storage and Disposal of CLAMENTIN F.C TABLET

-Storage

Store at temperature below 30°C. Protect from light and moisture.

-Disposal

CLAMENTIN F.C TABLET should be disposed when your doctor or pharmacist tells you to stop taking this tablet or it has passed the expiry date. Ask your doctor and pharmacist how to dispose medicines that are no longer required.

Product description

-What it looks like

CLAMENTIN F.C TABLET 375mg:

A white film coated tablet, one side impressed with "YSP4"



CLAMENTIN F.C TABLET 625mg:

An elliptical white film coated tablet, one side impressed with "YSP 3"



-Ingredients

-Active ingredients:

Amoxicillin Trihydrate
Clavulanate Potassium

-Inactive ingredients:

Cellulose
Magnesium Stearate
Silicon Dioxide
Sodium Starch Glycolate
Titanium Dioxide
Aerosil
Triethyl Citrate
Isopropyl Alcohol
Methylene Chloride

-MAL number

CLAMENTIN F.C TABLET 375mg:

MAL07124623AZ

CLAMENTIN F.C TABLET 625mg:

MAL07124630AZ

Manufacturer

Yung Shin Pharmaceutical Ind. Co., Ltd.
(Taichung Youth Factory III)
No. 26 Kung Liu Rd. Youth Industrial District, Taichia, Taichung, Taiwan, R.O.C

Product Registration Holder

Y.S.P Industries (M) Sdn. Bhd.
Lot 3,5 & 7, Jalan P/7, Section 13, Kawasan Perindustrian Bandar Baru Bangi, 43000 Kajang, Selangor Darul Ehsan, Malaysia.

Date of revision

21/4/2016

Serial Number

BPFK(R4/1)200416/00116