

# BAGES CREAM

Methyl salicylate/Menthol/Camphor (30 % w/w / 10 % w/w / 4 % w/w)

## What is in this leaflet

1. What Bages is used for
2. How Bages works
3. Before you use Bages
4. How to use Bages
5. While you are using it
6. Side effects
7. Storage and Disposal of Bages
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

## What Bages is used for

*Bages* is used for temporarily relieves of minor aches and pains of muscles and joints.

## How Bages works

Methyl salicylate, menthol and camphor are external analgesics which stimulate sensation of warmth and/or cold. This produces a counter-irritant response which provides temporary relief of minor aches and pains of muscles and joints associated with simple backache, arthritis, strains and sprains.

## Before you use Bages

### - When you must not use it

Do not use *Bages*:

- If you are allergic to any of the ingredients
- If your child is below 2 years old.

### *Pregnancy and lactation*

Do not take *Bages* if you are pregnant, trying to get pregnant or think you may be pregnant.

Do not take *Bages* if you are breast-feeding. Ask your doctor or pharmacist for advice before taking any medicine.

### - Before you start to use it

This product is for external use only. Do not apply to wounds or damaged skin or over large areas of the body or bandage.

### - Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop especially, "blood thinners" (e.g. warfarin).

## How to use Bages

### - How much to use

Follow all directions given to you by your doctor and pharmacist carefully. They may differ from the information contained in this leaflet.

If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

Apply to affected area 2-3 times daily.

### - When to use it

Use as directed by your doctor or pharmacist.

### - How long to use it

Continue taking *Bages* for as long as your doctor recommends.

If condition worsens, or if symptoms persist for more than 7 days or clear up and occur again within a few days, discontinue use of this product and consult a doctor.

### - If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it.

Apply the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

### - If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Symptoms include dizziness, noise or ringing in the ears, deafness, sweating, nausea and vomiting, headache and confusion, and may be controlled by reducing the dosage.

## While you are using it

### - Things you must do

Take your medicine exactly as your doctor has told you.

Tell all the doctors, dentists and pharmacists treating you that you are taking *Bages*.

Tell your doctor immediately if you become pregnant while taking this medication.

### - Things you must not do

Do not stop taking the medicine unless advised by your doctor.

Do not take any new medicines without consulting your doctor or pharmacist.

Do not give *Bages* to anyone else, even if they have the same symptoms or condition as you.

### - Things to be careful of

Caution should be exercised when older children are treated.

It is dangerous to place any camphor containing product into the nostril of children. A small amount applied this way may cause immediate collapse.

## Side effects

Like all medicines, *Bages* can cause side effects, although not everybody gets them.

Visit your doctor or pharmacist immediately if you experience any side effects after taking this medicine.

Side effects are skin irritation, redness or swelling, burning or stinging sensation. Dizziness or nausea may also occur.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website [np.ra.moh.gov.my](http://np.ra.moh.gov.my) (Public → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI))

## Storage and Disposal of Bages

### - Storage

Keep medicines out of reach of children.

Store at a temperature not more than 30°C.

# BAGES CREAM

Methyl salicylate/Menthol/Camphor (30 % w/w / 10 % w/w / 4 % w/w)

---

## Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will you protect the environment.

## **Product Description**

### - What it looks like?

White, smooth, viscous, non-greasy cream with characteristic and pungent odor

### - Ingredients

- Active ingredients:

Methyl salicylate, Menthol and Camphor

- Inactive ingredients:

Butylated hydroxytoluene, Atlas G 1726, Cyclogol NI wax, Lutrol F68, Sodium lauryl sulfate, Simethicone emulsion 30%, Germaben II and Purified water

### - MAL Number:

MAL06100732X

## **Manufacturer**

Unison Laboratories Co., Ltd.  
39 Moo 4, Klong Udomcholjorn,  
Muang Chachoengsao,  
Chachoengsao 24000 Thailand

## **Product Registration Holder**

Medispec (M) Sdn. Bhd.  
No. 55 & 57, Lorong Sempadan 2  
(Off Boundary Road) 11400 Ayer  
Itam, Penang, Malaysia

## **Date of Revision**

6.3.2018

## **Serial Number:**

NPRA(R4/1)180118/00018