

DIPYRIDAMOLE TABLET 75 mg®

DIPYRIDAMOLE TABLET 75 mg

What is in this leaflet:

1. What Dipyridamole Tablet 75 mg is used for
2. How Dipyridamole Tablet 75 mg works
3. Before you use Dipyridamole Tablet 75 mg
4. How to take Dipyridamole Tablet 75 mg
5. While you are using Dipyridamole Tablet 75 mg
6. Side Effects
7. How to store and dispose
8. Product description

What Dipyridamole Tablet 75 mg is used for

Dipyridamole Tablet 75 mg is being used to inhibit the aggregation of platelets, usually conjunction with warfarin, following cardiovascular or renal surgery.

How Dipyridamole Tablet 75 mg works

It contains dipyridamole. This belongs to a group of medicines called anti-platelet agents. Dipyridamole helps to stop blood clots forming, particularly when you have artificial heart valves.

Before you use Dipyridamole Tablet 75 mg

Concurrent use of Dipyridamole Tablet 75 mg and heparin may cause increased danger of hemorrhage.

Dipyridamole Tablet 75 mg should be used with great caution in patients having propensity toward hypotension.

Continuous therapy with Dipyridamole Tablet 75 mg may required for 2 or 3 months before its full therapeutic efficacy is reached in the prevention of angina.

How to take Dipyridamole Tablet 75 mg

How much to take

Always take Dipyridamole Tablet 75 mg exactly as your doctor has told you. If you are not sure, check with your doctor or pharmacist.

Doses:

Adults (including elderly): 300-600mg a day.

Children: dosage has not been established.

When to take it

Swallow the tablets with water before meals.

If you forget to take it

If you miss a dose of Dipyridamole Tablet 75 mg, take it as soon as possible. Unless if nearly time for next dose, skip the missed dose and go back to your regular dosing schedule. Do not take 2 doses at once.

If you take too much (overdose)

If you have taken more than the prescribed dose contact your doctor immediately. Signs of an overdose include headache, stomach upsets and low blood pressure.

While you are using Dipyridamole Tablet 75 mg

1. Do not take more than the recommended dose.
2. You are advised to avoid alcohol.
3. If you have been told that you have an intolerance to some sugars, contact your doctor before taking this medicine.

Side Effects

Side effects include dizziness, lightheadedness or fainting, flushing of the face, headache, nausea or vomiting, skin rash, gastrointestinal disturbances and weakness. These side effects may be more frequent with increase dosage.

If you get any of following side effects, stop taking Dipyridamole Tablet 75 mg and see your doctor as soon as possible: worsening of your symptoms of heart disease, faster heart rate, increased bleeding after an operation, red mark on the skin and bruising.

How to store and dispose

Storage

Keep container tightly closed in a dry place, below 25°C. Protect from light. Keep out of the reach of children.

Disposal

Do not dispose the medicine into wastewater or household garbage. Ask your pharmacist how to dispose of medicines no longer required.

Product Description

What it looks like?

Dipyridamole Tablet 75 mg is a round, reddish orange sugar coated tablet. The pack sizes are: Plastic Container of 30 tablets and Plastic Container of 500 and 1000 tablets (for Tender or Export only).

Ingredients

The active substance is dipyridamole. Each tablet contain is 75 mg of active ingredients.

The other ingredients are lactose, povidone, sodium starch glycolate, magnesium stearate, talc powder, acacia, calcium carbonate, maize starch, titanium dioxide, light magnesium carbonate, color FDC yellow No. 6, plain syrup, PVP shellac, and polishing solution.

Registration Numbers:

Dipyridamole Tablet 75 mg:
MAL19860181A

Manufactured by:

Idaman Pharma Manufacturing Sdn Bhd
(661901-P)

Lot 24, Bakar Arang Industrial Estate,
Sungai Petani, Kedah, Malaysia.

Date of revision:

Marketing Authorisation Holder

August 10th, 2011

IDAMAN PHARMA MANUFACTURING SDN BHD