ORALON DENTAL PASTE

Triamcinolone Acetonide (0.1% w/w)

What is in this leaflet:

- 1. What Oralon Dental Paste is used for
- 2. How Oralon Dental Paste works
- 3. Before you use Oralon Dental Paste
- 4. How to use Oralon Dental Paste
- 5. While you are using Oralon Dental Paste
- 6. Side Effects
- 7. Storage and Disposal of Oralon Dental Paste
- 8. Product Description
- 9. Manufacturer and Product Registration Holder
- 10. Date of Revision

What Oralon Dental Paste is used for

Oralon dental paste is indicated as adjunctive treatment and temporary relieving inflammation and ulceration inside the mouth that is not related to herpes infection.

How Oralon Dental Paste works

Triamcinolone acetonide is the active ingredient of Oralon Dental Paste. Triamcinolone acetonide is used for narrowing the blood vessels and relieve inflammation and itching on the skin or mucous membranes.

Before you use Oralon Dental Paste

-When you must not use it

- If you are allergic (hypersensitive) to triamcinolone acetonide or any of the other ingredients of Oralon Dental Paste.
- If you have herpes lesions of known viral origin.

-Before you start to use it

You must tell your doctor if you have following condition as Oralon Dental Paste should not be used in the following condition:

- i) presence of infection.
- ii) tuberculosis (bacteria infection that mainly affect the lung), stomach ulcer, or diabetes.

If there is no symptoms improvement occurred in 7 days after you use Oralon Dental Paste, you should contact your doctor as additional investigations into the condition of oral lesion should be conducted by your doctor.

Pregnancy and breastfeeding

You should talk with your doctor if you are pregnant, think that you may be pregnant, planning to become pregnant, or breast-feeding. Oralon Dental Paste should not be used in women of child-bearing potential and particularly during early pregnancy unless the potential benefits outweighed the possible hazards.

-Taking other medicines

Always inform your doctor and pharmacist if you are taking any other medicines, including herbal tonics, supplements and medicines that you buy without prescription.

How to use Oralon Dental Paste

-How much to use

- Use Oralon Dental Paste exactly as your doctor has told you. You should check with your doctor or pharmacist if you are not sure.
- A small dab of the Oralon Dental Paste is to applied to the lesion, two or three times a days.
- Applying the paste with a cotton buds will help to eliminate any possible absorption from contact with skin.
- The paste should be pressed, not rubbed, on the lesion. Rubbing the paste on the lesion will result in a granular, gritty sensation and cause it to crumble. A smooth, slippery film forms after application.

-When to use it

Oralon Dental Paste should be applied to the lesion after meals and at bed time.

-How long to use it

To apply for as long as your doctor recommends it.

-If you forget to use it

- If you forget a dose, apply it as soon as you remember it.
- However, if it is nearly time for the next dose, skip the missed dose.
- Do not apply a double dose to make up for a forgotten dose.

-If you use too much (Overdose)

Using too much Oralon Dental Paste may increase the chance of skin irritation such as burning, itching, blistering or peeling.

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

While you are using Oralon Dental Paste

-Things you must do

You should use Oralon Dental Paste exactly as your doctor has prescribed.

-Things you must not do

Do not give Oralon Dental Paste to anyone else, even if their symptoms seem similar or they have same condition as you. Your doctor has prescribed Oralon Dental Paste for you and your condition.

-Things to be careful of

If you experience any side effects, infection or irritation, you should consult your doctor for advice.

This medicine will reduce your oral immune system.

Side Effects

Like all medicines, Oralon Dental Paste can have side effects, although not everybody gets them. The following side effects may happen with this medicine:

- Signs of infection:
 Pain, redness or pus-containing blister.
- Signs of irritation:
 Burning, itching, blistering or peeling not present before therapy.

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

Consult your doctor or pharmacist if you are in doubt or for any further information.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website npra.moh.gov.my

(Public→Reporting Medicinal Problems/ Side Effects/ AEFI/ Vaccine Safety)

Storage and Disposal of Oralon Dental Paste

-Storage

Do not use after the expiry date stated on the pack. The expiry date refers to the last day of that month.

Oralon Dental Paste should be stored at temperature between 15 to 25 °C. The tube should be tightly closed after each usage.

Keep out of the reach and sight of children.

-Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

-What it looks like

A light brown paste in printed tubes with white caps. They are supplied in 5g of collapsible aluminium tubes.

- Ingredients

- Active Ingredient:

Triamcinolone Acetonide.

- Inactive ingredients:

Mineral Oil Heavy, White Petrolatum/ Soft Paraffin, White Wax, Cholesterol, Stearyl Alcohol, Gelatin, Pectin, Carboxymethylcellulose Sodium.

-MAL number:

MAL19920519AZ

Manufacturer and Product Registration Holder

Malaysian Pharmaceutical Industries Sdn. Bhd.

Plot 14, Lebuhraya Kampung Jawa, 11900 Bayan Lepas, Penang, Malaysia.

Date of Revision

04/05/2017

Serial number:

BPFK (R4/1) 030517/00091