

# PEACE TABLET

Triprolidine hydrochloride/ Pseudoephedrine hydrochloride (2.5mg/60mg)

## What is in this leaflet

1. What PEACE TABLET is used for
2. How PEACE TABLET works
3. Before you use PEACE TABLET
4. How to use PEACE TABLET
5. While you are using it
6. Side effects
7. Storage and Disposal of PEACE TABLET
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

## What PEACE TABLET is used for

PEACE TABLET is used for symptomatic relief of sneezing, stuffy nose and rhinorrhea caused by seasonal, allergic and vasomotor (nonallergic) rhinitis (inflammation of the mucous membrane inside the nose).

## How PEACE TABLET works

Triprolidine belongs to a class of medications known as antihistamine. It works by blocking the action of histamines, biological substances in the body that produces symptoms of sneezing, itching, watery eyes and runny nose.

Pseudoephedrine is a type of sympathomimetic agent that works by constricting blood vessels in the respiratory tract, leading to a reduction in mucosal edema (swelling) and nasal congestion.

## Before you use PEACE TABLET

### - When you must not use it

- You must not take PEACE TABLET if you have cardiovascular (heart or blood vessel) disease, endocrine (hormonal disorder) diseases and known hypersensitivity to this preparation.
- If you are under treatment with Monoamine Oxidase Inhibitors (MAOIs) and within 2 weeks of stopping it, you should avoid taking PEACE TABLET.

### - Before you start to use it

- Consult your doctor and pharmacist regarding the consumption of PEACE TABLET if you are pregnant or breastfeeding.
- Tell your doctor or pharmacist if you are allergic to any other medicines or you have any other health related problems such as narrow-angle glaucoma (iris of eyes is pulled or pushed forward), pyloroduodenal obstruction (obstruction on the gastric outlet), prostatic hypertrophy (non-cancerous enlargement of prostate gland), diabetes (high blood sugar), ischemic heart disease also known as coronary heart disease (narrowing of arteries), hyperthyroidism (overactivity of thyroid gland that leads to rapid heartbeat) and severe impairment of kidney and liver function.

### - Taking other medicines

Avoid concurrent use of PEACE TABLET with:

- Sympathomimetic agents (eg. Decongestant, tricyclic antidepressants, appetite suppressants, amphetamine-like psychostimulants or monoamine oxidase inhibitors)
- Antihypertensive agents (methyldopa, guanethidine)
- Furazolidone antibacterial agent.

Seek for your doctor or pharmacist advice if you are taking any of the medication mentioned or other medicines, including those you have bought without a prescription.

## How to use PEACE TABLET

### - How much to use

Adults and children >12 years: 1 tablet, 3-4 times daily

Children 6-12 years: ½ tablet, 3-4 times daily

### - When to use it

PEACE TABLET should be taken according to the time advised by your doctor or pharmacist. It should be taken orally with or without food.

### - How long to use it

PEACE TABLET should be taken regularly and as long as your doctor advised you to do so.

### - If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it.

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

### - If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Taking too many tablets may cause:

- drowsiness
- dizziness
- incoordination
- weakness
- convulsion
- respiratory depression
- tachycardia (a faster than normal heart rate).

## While you are using it

### - Things you must do

Take your medicine exactly as your doctor has told you.

Tell all the doctors, dentists, and pharmacists treating you that you are taking PEACE TABLET.

Tell your doctor immediately if you become pregnant while taking this medication.

- Things you must not do

Do not stop taking the medicine unless advised by your doctor.

Do not take any new medicines without consulting your doctor.

Do not give PEACE TABLET to anyone else, even if they have the same symptoms or condition as you.

- Things to be careful of

- Avoid alcohol consumption during the course of treatment with PEACE TABLET.
- Be careful if you drive or do anything that requires you to be alert and awake after consuming PEACE TABLET.
- It should be used with caution in children under 6 years old.

**Side effects**

PEACE TABLET may cause dryness of mouth, nose and throat, sedation, sleepiness, dizziness, disturbed co-ordination, hypotension (low blood pressure), headache, palpitation (irregular heartbeat), tachycardia, epigastric distress, anorexia, nausea, vomiting and thickening of bronchial secretions. Please tell your doctor or pharmacist if you experience any other side effects.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website [portal.bpfk.gov.my](http://portal.bpfk.gov.my) (Consumers →Reporting).

**Storage and Disposal of PEACE TABLET**

- Storage

Store below 30°C, protect from light & moisture. Keep in a tight container.


- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

**Product Description**

- What it looks like

PEACE TABLET is a light yellow square tablet, one side impressed with the 'YS', and the other side with the

mark "".

- Ingredients

- Active ingredients.  
Triprolidine hydrochloride  
pseudoephedrine hydrochloride
- Inactive ingredients  
Sodium Starch Glycolate  
Talc  
Magnesium Stearate  
Neusilin  
Avicel  
Isopropyl Alcohol  
Riboflavin

- MAL number:

MAL19970980AZ

**Manufacturer & Product Registration Holder**

Y.S.P Industries (M) Sdn. Bhd.  
Lot 3,5 & 7, Jalan P/7, Section 13,  
Kawasan Perindustrian Bandar Baru  
Bangi,  
43000 Kajang, Selangor Darul Ehsan,  
Malaysia.

**Date of revision**

26 October 2015

**Serial number:**

BPFK(R4/1)051015/00333