

TOLNADERM LOTION

Tolnaftate (1% w/v)

What is in this leaflet

1. What Tolnaderm Lotion is used for?
2. How Tolnaderm Lotion works?
3. Before you use Tolnaderm Lotion
4. How to use Tolnaderm Lotion
5. While you are using Tolnaderm Lotion
6. Side effects
7. Storage and Disposal of Tolnaderm Lotion
8. Product description
9. Manufacturer and Product Registration Holder
10. Date of revision

This leaflet answers some common questions about Tolnaderm Lotion. It does not contain all the available information. It does not take the place of talking to your pharmacist or doctor.

All medicines have risks and benefits. Your pharmacist or doctor has weighed the risks of you taking Tolnaderm Lotion against the benefits they expect it will have for you.

Keep this leaflet with the medicine.

You may need to read it again.

What Tolnaderm Lotion is used for

Tolnaderm Lotion is used to treat fungal skin infections such as body ringworm, ringworm of the face and neck, athlete's foot and jock itch.

How Tolnaderm Lotion works

Tolnaderm Lotion contains the active ingredients Tolnaftate. Every bottle contains Tolnaftate 1% w/v in Polyethylene glycol.

Tolnaftate is broad spectrum in the treatment of the skin in fungal infection.

Ask your pharmacist or doctor if you have any questions about this medicine.

Before you use Tolnaderm Lotion

-When you must not use it

Do not apply Tolnaderm Lotion if:

- You are allergic (hypersensitive) to Tolnaftate or to any of the other ingredients in this medicine.
- You are pregnant, breast-feeding or trying for a baby, tell your doctor or midwife before using Tolnaderm Lotion. If you have informed your doctor or midwife already, follow his/her instructions carefully.

-Before you start to use it

Tell your pharmacist or doctor:

- If you are allergic (hypersensitive) to Tolnaftate or to any of the other ingredients in this medicine.

Pregnancy and breast feeding:

- Safe use during pregnancy and lactation has not been established.
- Ask your doctor or pharmacist for advice before taking this medicine if you are pregnant or breast-feeding.

-Taking other medicines

Tell your pharmacist or doctor if you are taking any other products (such as over the counter drug, prescribed drug and herbal products).

How to use Tolnaderm Lotion

-How much to use

Do not use this medicine to children under 2 years of age. Consult with your doctor if you want to give this medicine to children under 2 years of age.

If Tolnaderm Lotion has been prescribed for you by your doctor, follow any instructions he/she may have given you.

If you purchased this product without a prescription, follow these directions closely:

- Wash and dry affected area. Apply small amount of lotion and spread evenly on infected

area 2 or 3 times daily for 2 to 3 weeks.

- Or 4 to 6 weeks if thickening of the skin has occurred.

-When to use it

2 or 3 times daily for 2 to 3 weeks after wash and dry affected area.

-How long to use it

The treatment should be continued for at least two weeks after the disappearance of all signs of infection to prevent re-infection. The symptoms of your infection, such as itching or soreness, should improve within a few days of treatment although signs such as redness and scaling may take longer to disappear. If symptoms persist, consult your doctor.

-If you forget to use it

Apply the lotion as soon as possible and then continue the rest of your treatment as usual.

-If you use too much (overdose)

No information is available concerning overdosage in human beings following topical application.

While you are using Tolnaderm Lotion

-Things you must do

You can help the treatment to work if you follow simple self-help tips:

- Although the infection area will itch, try do not scratch. Scratching will damage the surface of the skin and cause the infection to spread further
- Keep the affected skin areas clean.
- Pay particular attention to drying the skin, but avoid excessive rubbing.
- Do not share towels, bath mats, etc. with other people as you could spread the infection to them.
- Always wash your hands after treating the infection to prevent it from spreading.

TOLNADERM LOTION

Tolnaftate (1% w/v)

-Things you must not do

Tolnaderm Lotion is for external use only. Do not put the lotion in your mouth or swallow it. If the lotion is swallowed accidentally, tell your doctor straight away or contact the Accident and Emergency Department of your nearest hospital. If you accidentally get solution in your eyes or mouth, wash immediately with water and contact your doctor. Avoid contact with eyes and mucous membranes (a lubricating membrane lining an internal surface or an organ, as mouth, nose, anus and vagina).

-Things to be careful of

Stop using this medicine if you are allergic to any of the listed ingredients.

Side effects

Tolnaderm Lotion can cause side effects, although not everybody gets them. As with all medicines, some people may be allergic to the lotion. If you are allergic, a reaction will occur soon after you start using it. If you experience an allergic reaction, stop using Tolnaderm Lotion and tell your doctor straight away or contact the Accident and Emergency Department of your nearest hospital. Signs of an allergic reaction may include:

- Rash.
- Swallowing or breathing problems.
- Swelling of your lips, face, throat or tongue.
- Weakness, feeling dizzy or faint.
- Nausea.

After you apply the lotion you might experience:

- Itching, rash, blister, burning, discomfort, swelling, irritation or peeling of skin.

If you experience any of the above effects or react badly to the solution in any other way not listed in this leaflet, tell your doctor or pharmacist immediately.

You may report any side effects or adverse drug reactions directly to

the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website npra.moh.gov.my (Public → Reporting Medicinal Problem / Side Effects / AEFI / Vaccine Safety).

Storage and Disposal of Tolnaderm Lotion

-Storage

Store below 30°C. Keep container well closed. Keep in a cool dry place, protected from light. Keep out of reach of children.

-Disposal

Dispose the medicine after the expiry date. Consult your pharmacist on how to safely dispose of any medicines no longer required.

Product description

-What it looks like

Tolnaderm Lotion is available in dropper bottles containing 10ml of clear viscous liquid.

-Ingredients

-Active ingredient(s):
Tolnaftate 1% w/v.

-Inactive ingredients:
Butylated Hydroxytoluene,
Polyethylene Glycol 400.

-MAL number:

MAL19930440X

Manufacturer

HOE Pharmaceuticals Sdn. Bhd.
Lot 10, Jalan Sultan Mohamed 6,
Bandar Sultan Suleiman,
42000 Port Klang, Selangor,
Malaysia.

Product Registration Holder:

HOE Pharmaceuticals Sdn. Bhd.
Lot 10, Jalan Sultan Mohamed 6,
Bandar Sultan Suleiman,
42000 Port Klang, Selangor,
Malaysia.

Date of revision

03/07/2017

Serial Number. :

BPFK(R4/1)21062017/00144