MICOZOL CREAM

Miconazole nitrate (2% w/w)

What is in this leaflet

- 1. What Micozol is used for
- 2. How Micozol works
- 3. Before you use Micozol
- 4. How to use Micozol
- 5. While you are using Micozol
- 6. Side effects
- 7. Storage and Disposal of Micozol
- 8. Product description
- 9. Manufacturer and Product Registration holder

10.Date of revision

What Micozol is used for

It is used for treatment of fungal infections of the skin and nails, including candidiasis, tinea and pityriasis versicolor (fungal infection of the skin)

How Micozol works

Micozol destroys both the fungus that causes the infection and some of the associated bacteria which may also be present.

Before you use Micozol

- When you must not use it
- allergic to miconazole or other ingredients of Micozol.
- Before you start to use it

Avoid contact with eyes. Do not bandage or wrap the treated skin area unless directed to do so by physician. Do not use in children under 2 years of age except under the advice and supervision of a doctor. It irritation occurs or if there is no improvement within 4 weeks, discontinue use and consult a doctor or pharmacist. Safety for use in pregnancy and during breastfeeding has not been established. Inform your doctor if you are pregnant or breastfeeding.

- Taking other medicines

The medicine is for topical application and will usually have minimal effect on other medication. However, please inform the doctor or pharmacist if taking other medicines.

How to use Micozol

- How much to use

Skin infections - Apply sparingly and gently rub into the affected area once or twice daily until lesion has completely disappeared.

Nail infections - Apply sparingly to the affected nail twice daily; combine with oral antimycotic or with a non-porous dressing, if necessary. Medication to be continued for full course of treatment to reduce the possibility of recurrence for 2 weeks.

- When to use it

Please follow the advice of the doctor or pharmacist.

- How long to use it

You should check with your doctor or pharmacist.

- If you forget to use it

If you forget to use this medicine, use it as soon as you remember. Then continue using it at your usual time.

- If you use too much (overdose)

If you have used too much Micozol, contact your doctor, an emergency department or a poisoning center immediately.

While you are using Micozol

Things you must do

Inform your doctor of all the medications you are taking.

- Things you must not do

Do not stop using medication although you feel better after a few days, until the prescribed course is finished.

- Things to be careful of

If you are pregnant, become pregnant, or are breastfeeding, you should not use Micozol unless advised by your doctor.

Side effects

Local irritation and occasionally sensitivity may occur at the site of application. If any of the side effects become serious or any other side effects not listed in this leaflet, please inform the doctor or pharmacist. You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website npra.moh.gov.my (Public \rightarrow Reporting Medicinal Problems / Side Effects / AEFI / Vaccine Safety).

Storage and Disposal of Micozol

- Storage

Store in cool, dry place below 30°C. Protect from light. Keep out of reach of children.

- Disposal

Ask your pharmacist how to dispose of medicines no longer required.

Product description

What it looks like

White colour cream.

- Ingredients:

- Active ingredient: miconazole nitrate
- Inactive ingredients: cetomacrogol emulsifying wax, white soft paraffin, liquid paraffin, chlorocresol, sodium citrate, disodium edetate and purified water.

- <u>MAL number</u>:

MAL19900276AZ

Manufacturer and Product Registration Holder

Prime Pharmaceutical Sdn. Bhd. 1505, Lorong Perusahaan Utama 1 Taman Perindustrian Bukit Tengah 14000 Bukit Mertajam, Penang, Malaysia.

Date of revision

19/04/2017

Serial number

BPFK(R4/1)190417/00086