

BREACOL EXPECTORANT FOR ADULTS

Guaifenesin (100mg/5ml)

Side effects

Rarely, the syrup may cause gastrointestinal discomfort and drowsiness.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website npra.moh.gov.my (Public → Reporting Medicinal Problems / Side Effects / AEFI / Vaccine Safety).

Storage and Disposal of Breacol Expectorant for Adults

- Storage

- Do not use Breacol Expectorant for Adults after the expiry date which is stated on the pack after 'Exp'.
- Do not store Breacol Expectorant for Adults above 30°C.

- Disposal

- Please discard the bottle once medication is finished or expired.

Product description

- What it looks like

A thick amber coloured liquid, free from foreign matter. A menthol and apricot odour taste is experienced

- Ingredients

- Active ingredient: Guaifenesin
- Inactive ingredient: sodium benzoate, potassium sorbate, sodium citrate, menthol, citric acid anhydrous, Saccharin Sodium, liquid glucose, caramel liquid, carboxymethyl cellulose, apricot flavour and purified water

What is in this leaflet

1. What Breacol Expectorant for Adults is used for
2. How Breacol Expectorant for Adults works
3. Before you take Breacol Expectorant for Adults
4. How to take Breacol Expectorant for Adults
5. While you are taking Breacol Expectorant for Adults
6. Side effects
7. Storage and Disposal of Breacol Expectorant for Adults
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What Breacol Expectorant for Adults is used for

Breacol Expectorant for Adults is used as an expectorant for symptomatic relief of coughs due to colds and upper respiratory tract infections

How Breacol Expectorant for Adults works

Breacol Expectorant for Adults contains Guaifenesin which has expectorant action which facilitates the removal of mucous.

Before you take Breacol Expectorant for Adults

- When you must not use it

Known hypersensitivity to Guaifenesin or any other ingredients in the formulation"

- Before you start to take it

Talk to your doctor or pharmacist before taking this medicine.

Usage during pregnancy and lactation should be considered based on risk benefit.

Talk to your doctor or pharmacist if you are pregnant or breast-feeding.

- Taking other medicines

Please consult your doctor or pharmacist if you are taking other medicines.

How to take Breacol Expectorant for Adults

- How much to take

DOSAGE:

Adults: 12 years and above:

Take 2-4 teaspoons (10-20ml) every 4 hours as required.

- When to take it

Take Breacol Expectorant for Adults every 4 hours as required.

- How long to take it

If you are coughing for more than a week, please consult your doctor

- If you forget to take it

If you forget to take Breacol Expectorant for Adults, please take it at the next dose.

- If you take too much (overdose)

Excessive use may cause nausea and vomiting. If you take more than you should, tell your doctor or pharmacist.

While you are taking Breacol Expectorant for Adults

- Things you must do

Keep medicine out of reach for children.

- Things you must not do

Do not take more than the recommended dose.

- Things to be careful of

If you are about to have carcinoid syndrome tests, tell your doctor that you are taking this medicine. It may interfere with the results of these tests.

If symptoms persist, consult the doctor

- *MAL number:*

MAL13115228XC

Manufacturer:

Pharmaniaga Manufacturing Berhad
11A, Jalan P/1,
Kawasan Perusahaan Bangi,
43650 Bangi,
Selangor, Malaysia.

Product Registration Holder:

A.Menarini Singapore Pte. Ltd.
B-18-2, Level 18, The Ascent
Paradigm,
No.1, Jalan SS7/26A, Kelana Jaya,
47301 Petaling Jaya, Selangor Darul
Ehsan, Malaysia.

Date of revision:

23/05/2017

Serial Number:

BPFK(R4/1)230517/00109