

DYNAMOL TABLET

Paracetamol (500mg, 650 mg)

What is in this leaflet

1. What Dynamol Tablet is used for
2. How Dynamol Tablet works
3. Before you use Dynamol Tablet
4. How to use Dynamol Tablet
5. While you are using it
6. Side effects
7. Storage and disposal of Dynamol Tablet
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What Dynamol Tablet is used for

It is used for the relief of pain such as headache, nerve inflammation, muscle pain, toothache and fever.

How Dynamol Tablet works

Paracetamol acts by inhibiting the production of hormone-like substances that participate in a wide range of body functions such as the contraction and relaxation of muscle, the dilation and constriction of blood vessels, control of blood pressure and modulation of inflammation. It is considered to affect the brain that controls body temperature.

Before you use Dynamol Tablet

- When you must not use it

Tell your doctor if you hypersensitive to Paracetamol or other ingredients in this product.

This preparation contains PARACETAMOL. Do not take Paracetamol containing medicines at the same time.

- Before you start to use it

Pregnancy and lactation

No reports regarding adverse effects. Lactation: Amount too small to be harmful.

Ask your doctor or pharmacists for advice before taking Dynamol Tablet.

- Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

Paracetamol should not be taken concomitantly with

- Alcohol
- Cholestyramine (used to reduce cholesterol)
- Warfarin (prevent blood clot)
- Metoclopramide or Domperidone (treat nausea and vomiting)

How to use Dynamol Tablet

- How much to use

Dynamol 650 Tablet

Adults: 1 tablet

Children 7 to 12 years: ½ tablet

Dynamol Tablet 500mg

Adults: 2 tablets.

Children 7 to 12 years: ½ - 1 tablet.

The directions given may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

- When to use it

To be taken 3 to 4 times daily (Not more than 4 doses in any 24 hours unless prescribed by a physician). Use as directed by your doctor or pharmacist.

- How long to use it

Continue taking Dynamol Tablet for as long as your doctor recommends.

Do not exceed the recommended dose or use for more than 10 days unless prescribed.

- If you forget to use it

Take the missed dose as soon as you remember. If it almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

- If you use too much (overdose)

Features of Paracetamol poisoning are nausea and vomiting, followed by pain and tenderness in the right side of lower ribs area of the abdomen. Overdose can cause liver damage that is signified by yellowing of the skin and eyes, gastrointestinal bleeding and low blood glucose level.

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

While you are using it

- Things you must do

Take your medicine exactly as your doctor has told you.

Tell the doctor, dentist and pharmacist treating you that you are taking Dynamol Tablet.

- Things you must not do

Do not take any new medicines without consulting your doctor or pharmacist.

Do not give Dynamol Tablet to anyone else, even if they have the same symptoms or condition as you.

- Things to be careful of

This medicine should be used with care in patient with liver and kidney impairment.

Side effects

Side effects of the Dynamol Tablet usually mild. Nausea, gastro-intestinal upsets, rashes and allergic conditions have occurred.

Allergy alert: Paracetamol may cause severe skin reactions. Symptoms may include skin reddening, blisters or rash. These could be signs of a serious condition. If these reactions occur, stop use and seek medical assistance right away.

Visit your doctor or pharmacist immediately if you experience any side effects after taking this medicine. You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website nppra.moh.gov.my (Public → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI).

DYNAMOL TABLET

Paracetamol (500mg, 650 mg)

Storage and Disposal of Dynamol Tablet

- Storage

Keep out of reach of children
Protect from light
Keep the container tightly closed

Date of revision

11/04/2018

Serial Number:

NPRA(R4/1)040418/00092

Dynamol 650 Tablet

Store in dry place below 25°C

Dynamol Tablet 500mg

Store in dry place below 30°C

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- What it looks like

Dynamol 650 Tablet

Pink colour tablet, round, flat scored with 'DYNAMOL' marking

Dynamol Tablet 500mg

White colour tablet, round, flat scored with 'DYNAMOL' marking

- Ingredients

- Active ingredients

Paracetamol

- Inactive ingredients

Dynamol 650 Tablet

PVP-K 90, Carmoisine, Magnesium Stearate and Corn Starch

Dynamol Tablet 500mg

PVP-K 90, Magnesium Stearate, Lactose & Dried Starch

- MAL number

Dynamol 650 Tablet

MAL19990615XZ

Dynamol Tablet 500mg

MAL19910653XZ

Manufacturer & Product registration holder

Dynapharm (M) Sdn. Bhd.

2497, Mk 1, Lorong Perusahaan Baru 5,

Kawasan Perusahaan Perai 3, 13600

Perai, Pulau Pinang, Malaysia.