

PHARMANIAGA RIBOFLAVIN TABLET

Riboflavine (3mg)

What is in this leaflet

1. What Pharmaniaga Riboflavin Tablet is used for
2. How Pharmaniaga Riboflavin Tablet works
3. Before you take Pharmaniaga Riboflavin Tablet
4. How to take Pharmaniaga Riboflavin Tablet
5. While you are using Pharmaniaga Riboflavin Tablet
6. Side effects
7. Storage and disposal of Pharmaniaga Riboflavin Tablet
8. Product description
9. Manufacturer and Product Registration Holder
10. Date of Revision

What Pharmaniaga Riboflavin Tablet is used for

It is used in the treatment of riboflavine deficiency states.

How Pharmaniaga Riboflavin Tablet works

Riboflavin is required for the proper development and function of the skin, lining of the digestive tract, blood cells, and many other parts of the body.

Before you take Pharmaniaga Riboflavin Tablet

- When you must not to use it

If you have allergies to Riboflavine or any ingredients contained in this product.

- Before you start to take it

Tell your doctor or pharmacist if:

- You are pregnant or intend to become pregnant.
- You are breast-feeding or planning to breastfeed.

- Taking other medicines

Tell your doctor or pharmacist if you are taking any other medicines, including medicines that you buy without a prescription from a pharmacy, supermarket or health food shop especially:

- probenecid (used to prevent gout)
- tricyclic antidepressants (to treat depression)
- phenothiazines (to treat psychotic disorders).

How to use Pharmaniaga Riboflavin Tablet

Follow all directions given to you by your doctor or pharmacist carefully. They may differ from the information contained in this leaflet.

If you do not understand the instructions on the label of your medicine, ask your doctor or pharmacist.

- How much to use

Adults:

Treatment dose: Up to 30 mg daily.

Maintenance dose: 1 to 3 mg daily.

Children:

Treatment dose: 3 to 10mg daily

- When to use it

Use as directed by your doctor or pharmacist.

- How long to use it

Continue taking Pharmaniaga Riboflavin Tablet as long as your doctor recommends it.

- If you forget to use it

Take the missed dose as soon as you remember.

If it is almost time for your next dose, skip the dose you missed and take your next dose when you are meant to. Do not take a double dose to make up for the dose that you missed.

This may increase the chance of you getting unwanted side effects.

- If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

While you are using Pharmaniaga Riboflavin Tablet

- Things you must do

Take your medicine exactly as your doctor has told you.

Tell all the doctor, dentists and pharmacists treating you that you are taking Pharmaniaga Riboflavin

Tablet.

Tell your doctor immediately if you become pregnant while taking this medication.

- Things you must not do

Do not give Pharmaniaga Riboflavin Tablet to anyone else, even if their symptoms seem similar to yours.

Do not take your medicine to treat any other complaints unless your doctor or pharmacist tells you to.

- Things to be careful of

If you are pregnant, trying to get pregnant or are breast-feeding, you should not take Pharmaniaga Riboflavin Tablet without discussing it with your doctor first.

Side effects

Like all medicines, Pharmaniaga Riboflavin Tablet can cause side effects, although not everybody gets them. Riboflavine may cause yellow discoloration of urine. No cause for concern.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website npra.moh.gov.my (Public → Reporting Medicinal Problems / Side Effects / AEFI / Vaccine Safety).

Storage and disposal of Pharmaniaga Riboflavin Tablet

- Storage

Keep all medicines out of sight and reach of children.

Do not take this medicine after the expiry date stamped on the pack.

Store below 30°C. Protect from light.

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment

PHARMANIAGA RIBOFLAVIN TABLET

Riboflavine (3mg)

Product description

- What it looks like

Yellow, round, flat beveled edge tablet with double breakline and with marking KK on one side.

- Ingredients:

-Active ingredient(s):-

Riboflavine 3mg

-Inactive ingredient(s):-

Lactose, Magnesium Stearate,
Talcum, Starch, Kollidon

-MAL Number(s):-

MAL19961339X

Manufacturer:

Pharmaniaga Manufacturing Berhad
(60016-D),
11A, Jalan P/1,
Kawasan Perusahaan Bangi,
43650 Bandar Baru Bangi, Selangor
Darul Ehsan,
Malaysia.

Product Registration Holder:

Pharmaniaga Manufacturing Berhad
(60016-D),
11A, Jalan P/1,
Kawasan Perusahaan Bangi, 43650
Bandar Baru Bangi, Selangor Darul
Ehsan,
Malaysia.

Date of revision:

14/02/2018

Serial Number:

NPRA(R4/1)140218/00050