

# BEPROSALIC LOTION

Betamethasone Dipropionate / Salicylic Acid (0.064% w/w / 2% w/w)

---

## What is in this leaflet

1. What Beprosalic lotion is used for
2. How Beprosalic lotion works
3. Before you use Beprosalic lotion
4. How to use Beprosalic lotion
5. While you are using Beprosalic lotion
6. Side effects
7. Storage and Disposal of Beprosalic lotion
8. Product description
9. Manufacturer and Product Registration Holder
10. Date of Revision

This leaflet answers some common questions about Beprosalic lotion. It does not contain all the available information. It does not take the place of talking to your pharmacist or doctor.

All medicines have risks and benefits. Your pharmacist or doctor has weighed the risks of you taking Beprosalic lotion against the benefits they expect it will have for you.

## Keep this leaflet with the medicine.

You may need to read it again.

## What BEPROSALIC LOTION is used for?

BEPROSALIC LOTION is used for the relief of the inflammatory manifestations of corticosteroid-responsive dryness and thickening of the outer layer of the skin disorder which include psoriasis, chronic atopic dermatitis, neurodermatitis and seborrhoeic dermatitis.

## How BEPROSALIC LOTION works?

BEPROSALIC LOTION contains medicine called Betamethasone Dipropionate and Salicylic Acid. Betamethasone Dipropionate is a topically active fluorinated corticosteroid which has anti-inflammatory, anti-itching and blood vessel narrowing actions. Salicylic acid has keratolytic (ability to soften the top layer scales of the skin), antibacterial and antifungal actions.

**Ask your pharmacist or doctor if you have any questions about this medicine.**

## Before you use BEPROSALIC LOTION

### -When you must not use it

### **Do not take BEPROSALIC LOTION if:**

- You are allergic (hypersensitive) to Betamethasone Dipropionate, other corticosteroids, Salicylic acid, or other salicylates, or any components in the base.

Topical corticosteroids are not recommended in most viral infections of the skin and also tuberculosis (bacterial infection that mainly affects the lungs) and acne.

### -Before you start to use it

Tell your pharmacist or doctor:

- If you have infection on your skin and specify on the infected area.
- If you are using any other skin products.

### **Pregnancy and breast feeding:**

Ask your doctor or pharmacist for advice before taking this medicine if you are pregnant or breast-feeding.

It should be used only if the potential benefit justifies the potential risk to the fetus or nursing infant.

### -Taking other medicines

**Tell your pharmacist or doctor if you are taking any other products (such as over the counter drug, prescribed medicine and herbal products).**

## How to use BEPROSALIC LOTION

Always use BEPROSALIC LOTION exactly as your doctor has told you.

### -How much to use

Apply a few drops of BEPROSALIC LOTION to the affected areas twice daily or as directed by your doctor or pharmacist.

### -When to use it

Apply on the affected area once dry and clean.

### -How long to use it

Consult with your doctor or pharmacist for the treatment period.

### -If you forget to use it

If you forget to use your lotion at the right time, use it as soon as you remember, then carry on as before. Do not double the dose to catch up.

### -If you use too much (overdose)

Corticosteroids applied to the skin can be adsorbed in sufficient amounts to produce systemic effects such as adrenal suppression, manifestation of Cushing syndrome,

high blood sugar level and presence of sugar in the urine.

Salicylic acid in preparation applied to the skin can also be absorbed in sufficient amounts to produce salicylism such as nausea, vomiting and hear ringing in the ears.

If you suspect that you are overdosed, please consult your pharmacist or doctor. Bring any remaining medicine with you.

**While you are using  
BEPROSALIC LOTION**

-Things you must do

If you are about to start taking on any other new medicine, tell your pharmacist or doctor that you are taking BEPROSALIC LOTION.

-Things you must not do

Do not take more than the recommended dose unless your doctor tells you to.

Do not use in the eyes.

-Things to be careful of

Discontinue use if irritation sensitivity, excessive dryness or reaction develop.

Suitable precautions should be taken during prolonged treatment, when treating extensive body surface areas, when using the occlusive technique and when treating children (because of a larger skin surface area to bodyweight ratio).

Suitable precautions must be taken when treating stasis skin inflammation and other skin diseases with impaired circulation.

**Side effects**

Like all medicines, BEPROSALIC LOTION can cause side effects, although not everybody gets them.

Reported local adverse reactions are burning, itching, irritation, dryness, hair follicle inflammation, abnormal hair growth, acne eruptions, lightening of the skin, softening of the skin color, secondary infection, skin thinning, stretch mark and heat rashes. Salicylic acid preparations may cause skin inflammation.

If you get any side effects, even those not mentioned in this leaflet, tell your doctor or pharmacist.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website [portal.bpfk.gov.my](http://portal.bpfk.gov.my) (Consumers → Reporting)

**Storage and Disposal of  
BEPROSALIC LOTION**

-Storage

Store below 30°C.  
Keep container well closed.  
Protect from strong light.  
Keep out of reach of children.

-Disposal

Dispose the medicine after the expiry date.  
Consult your pharmacist on how to safely dispose of any medicines no longer required.

**Product description**

-What it looks like

BEPROSALIC LOTION is a colourless and slightly viscous solution. It is available in bottle containing 30ml.

-Ingredients

Active ingredients:  
Betamethasone Dipropionate  
0.064%w/w and Salicylic Acid  
2% w/w

Inactive ingredients:  
Methylparaben, Isopropyl Alcohol,  
Sodium Hydroxide and Purified  
Water.

-MAL number: MAL20021135A

**Manufacturer**

HOE Pharmaceuticals Sdn. Bhd.  
Lot 10, Jalan Sultan Mohd. 6,  
Bandar Sultan Suleiman,  
42000 Port Klang, Selangor,  
Malaysia.

**Product Registration Holder**

HOE Pharmaceuticals Sdn. Bhd.  
Lot 10, Jalan Sultan Mohd. 6,  
Bandar Sultan Suleiman,  
42000 Port Klang, Selangor,  
Malaysia.

**Date of revision**

17/02/2016

**Serial Number:**

BPFK(R4/1)160216/00040