

UPHAMOL 250 SYRUP (RED)

Paracetamol (250mg/5ml)

What is in the leaflet

1. What Uphamol Syrup is used for
2. How Uphamol Syrup works
3. Before you use Uphamol Syrup
4. How to use Uphamol Syrup
5. While you are using it
6. Side effects
7. Storage and Disposal of Uphamol Syrup
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of Revision

What Uphamol Syrup is used for

Uphamol Syrup is used for relief of pain and fever.

How Uphamol Syrup works

Uphamol Syrup contains paracetamol as an active ingredient. Paracetamol works by relieving pain and reducing high temperature.

Before you use Uphamol Syrup

- When you must not take it

Do not take this medicine if your child has an allergy to:

- paracetamol
- any of the other ingredients listed at the end of this leaflet

Allergy alert: Paracetamol may cause severe skin reactions. Symptoms may include skin reddening, blister or rash.

These could be signs of serious conditions. If these reactions occur, stop use and seek medical assistance right away.

Do not take Uphamol Syrup if your child is taking other products containing paracetamol.

Do not take this medicine after the expiry date printed on the pack or if the packaging is torn or shows signs of tampering. If it has expired or is damaged, return it to the pharmacist for disposal.

If you are not sure whether your child should start taking this medicine, talk to your doctor.

- Before you start to use it

Tell your doctor if your child has allergies to any other medicines, foods, preservatives or dyes.

Tell your doctor if you have or have ever had any other health problems / medical conditions, including:

- liver disease
- alcohol addiction problem

If your symptoms persist, please consult your doctor or pharmacist

- Taking other medicines

Tell your doctor or pharmacist if your child is taking any other medicines, including any that you get without a prescription from your pharmacy, supermarket or health food shop.

Some medicines and Uphamol Syrup may interfere with each other.

Your doctor and pharmacist have more information on medicines to be careful with or avoid while taking this medicine.

How to use Uphamol Syrup

Follow all directions given to you by your doctor and pharmacist carefully.

These directions may differ from the information contained in this leaflet.

If you do not understand the instructions on the pack, ask your doctor or pharmacist for help.

- How much to use

Children up to 1 year:
Take 60 – 120 mg

Children 1 to 5 years:
Take 125 – 250 mg

Children 6 to 12 years:
Take 250 – 500 mg

- When to use it

Take Uphamol Syrup every four to six hours or as your doctor has prescribed.

Do not take more than 4 times daily and never exceed the recommended dose unless prescribed by your doctor or pharmacist.

- How long to use it

Continue taking Uphamol Syrup for as long as the doctor recommends.

- If you forget to use it

Take the dose as soon as you remember, and continue to take it as you would normally.

If it is almost time for the next dose, skip the dose you missed and take the next dose when you are meant to.

Do not take a double dose to make up for the dose that you missed. This may increase the chance of getting an unwanted side effect.

- If you use too much (overdose)

Immediately telephone your doctor for advice or go to Accident and Emergency (Casualty) at your nearest hospital if you think that your child or anyone else may have taken too much Uphamol Syrup.

Do this even if there are no signs of discomfort or poisoning. Your child may need urgent medical attention.

While you are using it

- Things you must do

Tell all the doctors, dentists and pharmacists who are treating your child that he/she is taking Uphamol Syrup.

If your child is about to start taking any new medicine tell the doctor and pharmacist that your child is taking Uphamol Syrup.

- Things you must not do

Do not take Uphamol Syrup to treat any other complaints unless your doctor tells you to.

Do not give the medicine to anyone else, even if they have the same condition as your child.

- Things to be careful of

Side effects

All medicines can have side effects. Sometimes they are serious, most of the time they are not.

Ask for the advice of your doctor or pharmacist if you have any concerns about the effects of taking this medicine.

Liver damage may occur in prolonged use or overdose. Skin hypersensitivity reactions including skin rashes, swelling beneath the skin, Stevens Johnson Syndrome/Toxic Epidermal Necrolysis have been reported.

Tell your doctor or pharmacist if you notice anything else that is making your child feels unwell.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Monitoring by calling Tel: 03-78835550, or visiting the website npra.moh.gov.my (Public → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI).

Storage and Disposal of Uphamol Syrup

- Storage

Keep the medicine in its original packaging until it is time to take them. Keep container tightly closed.

Keep the medicine in a dry place where the temperature will stay below 30°C. Protect from light.

Do not store the medicine, or any other medicine, in the bathroom or near a sink.

Keep it where children cannot reach it.

- Disposal

If your doctor or pharmacist tells your child to stop taking this medicine or it has passed its expiry date, ask your pharmacist what to do with any medicine that is left over.

Product Description

- What it looks like

A red, orange flavoured syrup.

- Ingredients

- Active ingredient:
Each 5ml of syrup contains 250mg paracetamol.

- Inactive ingredients:

- Glycerine
- Methylparaben
- Propylparaben
- Carmoisine
- Saccharin Sodium
- Sugar
- Propylene Glycol
- Sweet Orange
- Chloroform Spirit

- MAL number

MAL19912688XZ

Manufacturer

Upha Pharmaceutical Mfg (M) Sdn Bhd (53667-M)
Lot 2&4, Jalan P/7, Section 13, Bangi Industrial Estate, 43650 Bandar Baru Bangi, Selangor Darul Ehsan

Product Registration Holder

Upha Pharmaceutical Mfg (M) Sdn Bhd (53667-M)
Lot 2&4, Jalan P/7, Section 13, Bangi Industrial Estate, 43650 Bandar Baru Bangi, Selangor Darul Ehsan

Date of Revision

15/05/2018

Serial Number:

NPRA (R4/1)080518/00143