DYNA B6 TABLET

Pyridoxine hydrochloride (100 mg)

What is in this leaflet

- 1. What Dyna B6 Tablet is used for
- 2. How Dyna B6 Tablet works
- 3. Before you use Dyna B6 Tablet
- 4. How to use Dyna B6 Tablet
- 5. While you are using it
- 6. Side effects
- 7. Storage and disposal of Dyna B6 Tablet
- 8. Product Description
- 9. Manufacturer and Product Registration Holder
- 10. Date of revision

What Dyna B6 Tablet is used for

Indicated for the treatment and prevention of pyridoxine deficiency states.

How Dyna B6 Tablet works

Vitamin B6 or Pyridoxine plays an important role in the body. It is needed to maintain the health of nerves, skin, and red blood cells.

Before you use Dyna B6 Tablet

- When you must not use it
 Do not use if you have any of the
 conditions:
- Hypersensitivity to the pyridoxine or other ingredients in this product.
- You have known inflammation of eye's optic nerve.
- <u>Before you start to use it</u>

 Pregnancy and lactation

 No reports regarding adverse effects.

Ask your doctor or pharmacists for advice before taking Dyna B6 Tablet.

- Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop expecially levodopa (used for Parkinson disease).

How to use Dyna B6 Tablet

- How much to use

For treatment and prevention of pyridoxine deficiency:

Adults, 1 to 4 tablets daily in divided doses.

For treatment of peripheral nerve inflammation caused by isoniazid therapy:

Adults, 1 to 2 tablets daily.

The directions given may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

- When to use it

Use as directed by your doctor or pharmacist.

- How long to use it

Continue taking Dyna B6 Tablet for as long as your doctor recommends.

- If you forget to use it

Take the missed dose as soon as you remember. If it almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

- *If you use too much (overdose)*No reports regarding overdose

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

While you are using it

- Things you must do

Take your medicine exactly as your doctor has told you.

Tell the doctor, dentist and pharmacist treating you that you are taking Dyna B6 Tablet.

- Things you must not do

Do not take any new medicines without consulting your doctor or pharmacist.

Do not give Dyna B6 Tablet to anyone else, even if they have the same symptoms or condition as you.

Side effects

Pyridoxine does not cause any side effects if taken according to the recommended dosage. Long term administration of large doses of pyridoxine is, however, associated with the development of severe peripheral nerve problem such as nerve pain, numbness and weakness.

Visit your doctor or pharmacist immediately if you experience any side effects after taking this medicine. You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website npra.moh.gov.my (Public → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI).

Storage and Disposal of Dyna B6 Tablet

- Storage

Keep out of reach of children
Protect from light
Store in dry place below 25°C
Keep the container tightly closed

- <u>Disposal</u>

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- <u>What it looks like</u>

Round, white colour tablets and biconvex

- Ingredients
- Active ingredients

Pyridoxine hydrochloride

DYNA B6 TABLET

Pyridoxine hydrochloride (100 mg)

- Inactive ingredients Lactose 200 Mesh, Magnesium Stearate, Corn Starch, PVP K-90
- <u>MAL number</u> MAL19940414XZ

Manufacturer and Product registration holder Dynapharm (M) Sdn. Bhd. 2497, Mk 1, Lorong Perusahaan Baru 5, Kawasan Perusahaan Perai 3, 13600 Perai, Pulau Pinang, Malaysia.

Date of revision 24/04/2018

Serial Number:

NPRA(R4/1)240418/00118