

BETACYCLIC OINTMENT

Betamethasone Dipropionate/Salicylic Acid (0.643mg/1g / 30mg/1g)

What is in this leaflet

1. What Betacylic Ointment is used for
2. How Betacylic Ointment works
3. Before you use Betacylic Ointment
4. How to use Betacylic Ointment
5. While you are using it
6. Side effects
7. Storage and Disposal of Betacylic Ointment
8. Product description
9. Manufacturer and Product Registration Holder
10. Date of revision

-How long to use it

Use as directed by your doctor or pharmacist.

-If you forget to use it

If you forget to Betacylic Ointment, apply it as soon as possible and then continue the rest of the treatment as usual.

-If you use too much (overdose)

Prolonged or excessive use of Betacylic Ointment might cause hypercorticism symptoms such as skin thinning. Growth retardation have been reported in children.

What Betacylic Ointment is used for

Relief of the inflammatory manifestation of thickening of the outermost layer of the skin and dry skin diseases that responds to corticosteroids such as psoriasis, dermatitis, lichen planus, eczema and other ichthyotic (coarse, dry, and scaly) conditions.

Excessive or prolonged use of topical preparations containing salicylic acid may cause symptoms of salicylism such as nausea, vomiting and hear ringing in the ears.

How Betacylic Ointment works

Betamethasone dipropionate has anti-inflammatory, antipruritic (anti-itching) and vasoconstrictive (ability to constrict blood vessel) actions. Salicylic acid has keratolytic (ability to soften the top layer scales of the skin), antibacterial and antifungal actions.

Seek medical attention immediately if found these symptoms.

While you are using it

-Things you must do

Always use it exactly as described on the label or as your doctor or pharmacist has told you. Promptly discontinue application of Betacylic Ointment if found sensitivity irritation or excessive dryness after using it.

Before you use Betacylic Ointment

-When you must not use it

If you are allergic to any components in Betacylic Ointment.

-Things you must not do

Do not contact it with your eyes and mucous membrane and keep it at the place that can be reached by children.

-Before you start to use it

Tell your doctor or pharmacist:

- if you have infection
- if you are pregnant or breast feeding
- if you are using other medicine.

-Things to be careful of

Precautions need to be taken if this cream is to be used for long-term, or in infants and children. Avoid extensive body use or occlusive technique when using Betacylic Ointment.

Betacylic Ointment should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

-Taking other medicines

Avoid using other medications on the areas you treat with Betacylic Ointment unless your doctor or pharmacist tells you to.

How to use Betacylic Ointment

-How much to use

Apply a thin film to the affected skin area twice daily.

-When to use it

In the morning and at night.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website portal.bpfk.gov.my (Consumers→Reporting)

Storage and Disposal of Betacylic Ointment

-Storage

Keep in a tight container. Store at temperature below 30°C. Protect from light and moisture.

-Disposal

Talk to your doctor or pharmacist about the proper disposal.

Product description

-What it looks like

Slight transparent, off white color ointment.

-Ingredients

Active Ingredient-
Betamethasone dipropionate
Salicylic acid

Inactive Ingredient-
Mineral oil and white petrolatum

-MAL number.

MAL 12025054A

Manufacturer

Y.S.P. Industries (M) Sdn. Bhd.
Lot 3,5 & 7, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru
Bangi, 43000 Kajang, Selangor D.E.,
Malaysia.

Product Registration Holder

Y.S.P Industries (M) Sdn. Bhd.
Lot 3,5 & 7, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru
Bangi,
43000 Kajang, Selangor Darul Ehsan
Malaysia.

Date of revision:

04/11/2015

Serial Number:

BPFK(R4/1)031115/00395